




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C07D 311/92, A61K 31/35		A1	(11) International Publication Number: WO 96/33988
			(43) International Publication Date: 31 October 1996 (31.10.96)
(21) International Application Number: PCT/US96/05673 (22) International Filing Date: 23 April 1996 (23.04.96) (30) Priority Data: 08/428,574 25 April 1995 (25.04.95) US (71) Applicant: WISCONSIN ALUMNI RESEARCH FOUNDATION [US/US]; 614 Walnut Street, Madison, WI 53707-7365 (US). (71)(72) Applicants and Inventors: BOOTHMAN, David, A. [US/US]; 7925 West Oakbrook Circle, Madison, WI 53717 (US). FRYDMAN, Benjamin, J. [US/US]; 402 Eau Claire No. 314, Madison, WI 53707 (US). WITIAK, Donald, T. [US/US]; 3046 Bosshard Drive, Madison, WI 53711 (US). (74) Agent: SARA, Charles, S.; DeWitt, Ross & Stevens, S.C., 8000 Excelsior Drive, Madison, WI 53717-1914 (US).		(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>With amended claims.</i>	

(54) Title: NOVEL SYNTHESIS AND USE OF BETA-LAPACHONE ANALOGS



I: Supercoiled p36B4 Substrate, II: Open-circular p36B4

(57) Abstract

3-Substituted- β -lapachone analogs and their use either alone or to augment chemotherapy or radiotherapy to induce programmed neoplastic cell death without exhibiting toxicity to surrounding normal cells are disclosed. In particular, 3-allyl- β -lapachones, 3-alkyl- β -lapachones and 3-halo- β -lapachones were found to be Topoisomerase (Topo I) inhibitors. When these analogs are used alone there is a reversible single-strand break in the DNA of neoplastic cells causing apoptosis and cell death in some cells. However, when these analogs are combined with chemotherapy or X-irradiation, an irreversible Topo I-mediated break is achieved. A new and more efficient chemical synthesis of the compounds is also disclosed.

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

NOVEL SYNTHESIS AND USE OF BETA-LAPACHONE ANALOGS

Background of the Invention

X-irradiation and chemotherapy are used to eliminate neoplastic cells. It has been found that a chemical post-treatment with β -lapachone analogs after X-irradiation greatly enhanced the lethality of the radiation and/or chemotherapy and destroy more neoplastic cells. Specifically, the β -lapachone post-treatment irreversibly damages the DNA of the neoplastic cells.

DNA is structurally damaged in various ways by different chemical and physical agents. For example, damage following ionizing radiation includes single and double-strand DNA breaks, apurinic/apyrimidinic alkaline-labile sites, altered basis, and intra- and inter-strand DNA protein cross-links. The predominant DNA lesions occurring after X-irradiation are single and double-strand breaks, and the production of the latter lesions correlate well with cytotoxicity. Each type of DNA lesion may stimulate different cellular responses which activate or induce a variety of DNA repair pathways. Along with the repair stimulation following radiation, mammalian cells respond to damage by altering gene expression and inducing post-translational modifications of proteins, including phosphorylation and ADP-ribosylation.

If damaged DNA regions are not repaired by the cell's enzymatic systems, the results are lethal or cause genetic rearrangements, ultimately leading to cell transformation and cancer. Rapid and accurate repair systems are important not only for cell survival, but for preventing genetic alterations which may lead to carcinogenesis. These facts are corroborated by a variety of human cancer-prone diseases, which are thought to result from deficiencies in DNA repair. These groups of diseases include the xeroderma pigmentosum, ataxia telangiectasia (whose cells are hypersensitive to X-irradiation) and Bloom's syndrome.

Conversely, if DNA repair is deliberately blocked or altered via administration of inhibitors, the lethal effects of anti-neoplastic agents can be greatly enhanced, often without creating carcinogenic consequences. The possibility of improving chemotherapy by inhibiting DNA repair has been suggested and tested years ago. In these prior studies, the major DNA repair inhibitor was caffeine, which was proposed to inhibit post-replication repair. Although the results of these studies were

promising, the extremely high concentrations of caffeine required to enhance radiotherapy or chemotherapy precluded their clinical utility.

Another class of DNA repair inhibitors proposed to enhance cancer chemotherapy were nicotinamide analogs, such as 3-aminobenzamide (3-ABA).
5 These compounds were initially proposed to be ligase inhibitors, but were later found to be inhibitors of poly (ADP-ribose) transferase (ADPRT). These compounds, therefore, block ADP-ribosylation which would otherwise stimulate DNA repair complexes and auto-down-regulate ADPRT itself. DNA repair inhibition by 3-ABA was based on this activity. However, 3-ABA, along with other nicotinamide
10 analogs, have not demonstrated significant clinical utility due to the fact that effective concentrations cannot be achieved.

β -lapachone is also a known inhibitor of cell survival after radiation because it is an inhibitor of DNA repair and Topoisomerase I. However, due to its extreme toxicity, this drug has not gained widespread use.

15 Camptothecin is also a well-known DNA-repair inhibitor. However, like β -lapachone, camptothecin and its derivatives are relatively toxic, causing lethality in healthy cells surrounding neoplastic sites. Consequently, many camptothecin derivatives of this compound have been synthesized to increase compound stability and water solubility. Camptothecin derivatives, as well as camptothecin itself, are
20 not related to β -lapachone or its derivatives and inhibits Topoisomerase I by different mechanisms than β -lapachone. Also, camptothecin and its derivatives are hydrophobic, making them poor candidates for pharmaceutical or medical applications.

Until now, little has been done to exploit DNA repair pathways for
25 chemotherapy or radiotherapy. Potential clinical utility exists for the development of drugs that modulate DNA repair enzymes and thereby enhance the lethality of DNA damaging agents. Novobiocin, an inhibitor of Topoisomerase II (Topo II), was useful as an adjunct to DNA damaging drugs. Pentoxiphylline, a caffeine analog, has been tested in phase I clinical trials and recent studies indicate that this compound, caffeine
30 and other methylxanthines, blocks DNA repair by allowing damaged cells to prematurely enter mitosis. These drugs prevent G₂ arrest in damaged cells, a stage which presumably allows DNA repair before cell division.

Past data indicate that Topoisomerase I (Topo I) is somehow involved in DNA repair and is affected by the aforementioned DNA repair inhibitors and, therefore, represents a prime target for radio- or chemotherapy. Areas of the mammalian genome undergoing high rates of transcription have higher Topo I levels and repair faster. Topo I is an essential enzyme required for transcription and replication. Topo I unwinds supercoiled DNA by breaking and rejoining one DNA strand as the enzyme rotates about its axis and passes the other DNA strand through the break. Topo I thereby relieves physical stress in DNA caused by RNA transcription and DNA replication. Unlike Topo II, Topo I levels within the cell do not vary as cells progress through their replicative cycle. Topo I is present in equivalent levels in log-phase as well as in quiescent human cells, which makes this enzyme an ideal target for chemo- and/or radio-therapeutic approaches. Many current chemical therapeutic regimens are directed against logarithmically growing cells (i.e. cells in S-phase). If radio- and chemo-therapeutic regimens can be developed that act upon non-cell cycle regulated enzymatic targets, resting as well as cycling tumor cells may be eliminated. This would aid in reducing resistance caused by recruitment of tumor cells from resting or hypoxic states.


Recent data indicates that Topo I inhibitors can cause a Topo I-mediated lesion modification of DNA in X-irradiated human cancer cells which were exposed to ionizing radiation. This Topo I mediated conversion of single-strand DNA breaks to double-strand breaks explains the radiation and chemotherapy synergy caused by β -lapachone derivatives. When Topo I binds to a plasmid containing a unique nick, the enzyme causes linearization of that substrate at an 800-fold greater rate compared to the enzyme's normal unwinding activity on intact DNA. Single-strand DNA regions or breaks are initially created by X-irradiation. Considerable evidence has accumulated to indicate that both mammalian and bacterial Topo I enzymes bind very avidly to single-strand DNA sequences and neither camptothecin nor β -lapachone affect such initial enzyme binding. Topo I has a very high affinity for nicked or single-stranded gapped, double strand DNA. After non-specifically binding to nicked DNA, the enzyme cuts the DNA strand opposite the nick resulting in linearization of the DNA substrate. The rate of this reaction was far more rapid (increased by a factor of 800 to 1,000) than nicking/unwinding activities upon undamaged plasmid

DNA. Thus, conversion of single-strand nicks into double-strand breaks by Topo I appears to be an extremely rapid and efficient reaction compared to the enzyme's actions upon intact undamaged double-stranded DNA.


The present hypothesis is that two pathways can exist for Topo I activity following radiation (see "Pathway 1" below).

PATHWAY 1

NORMAL PATHWAY



DRUG-MEDIATED SUICIDE PATHWAY



First, a normal pathway in which down-regulation of the enzyme is required for DNA repair. The second pathway involves a β -lapachone-analog-enzyme mediated suicide pathway in which a Topo I modulator somehow prevents the enzyme's down-regulation and/or changes the enzyme's activity site such that it can still bind, create a double-strand DNA break from a single-strand break and cause covalent attachment of the enzyme to the DNA break in spite of being modified. This results in a synergistic cell-killing mechanism between β -lapachone derivatives and radiotherapy or chemotherapy. Recent data indicate that cells undergoing apoptosis (or programmed cell death) may irreversibly degrade poly (ADP) ribosyl transferase (ADPRT), a key enzyme required for DNA repair and for controlling (i.e. down-regulating) Topo I. Thus, the combination of loss of ADPRT and subsequent uncontrolled Topo I activity may contribute to cell death, possibly via apoptosis. Recent data indicate that all three substituted β -lapachone derivatives can cause apoptosis indicating the use of these compounds alone as chemotherapeutic agents; in particular, these agents should be effective against prostate, colon, lung, melanoma, and breast cancers due to the elevated Topo I enzyme levels in these tumors compared to normal cells.

After cells receive ionizing radiation, it appears that the DNA repair complex, which probably contains DNA ligase, DNA polymerase- β , and ADPRT, competes with Topo I and other proteins for available single-strand DNA damaged sites. Creation of double-strand DNA lesions from single-strand DNA breaks has been reported under normal repair conditions and may be the result of the cell's inability to efficiently down-regulate Topo I. Normally, after ionizing radiation, ADPRT/DNA binding and subsequent enzymatic activities increase. ADPRT acts to stabilize the break (ADPRT/DNA binding) and its ADP-ribosylation activity, which increases due to DNA binding, can stimulate DNA repair and down-regulate Topo I. This combined action by ADPRT functions to stabilize DNA lesions, out-competes Topo I for these sites, and prevents conversion of single-stranded breaks to double-stranded breaks by this down-regulation of Topo I. The temporary down-regulation of Topo I is needed, therefore, to repair single-strand breaks. After some time (approximately 60 minutes in human cells), Topo I activity recovers and its unwinding activity may be needed to repair other types of more persistent DNA damage, such as DNA

protein cross-links, which can be induced by chemotherapeutic agents. In fact, Topo I inhibitors (such as camptothecin) have been used to augment cis-platin and melphalan cytotoxicities.

5 In the presence of a modulator of Topo I, such as the present invention of β -lapachone analogs, the processes of DNA repair is altered and DNA lesion modification by Topo I occurs. Somehow Topo I is no longer down-regulated by cellular increases in ADPRT in the presence of β -lapachone derivatives. The manner by which these compounds either prevent Topo I down-regulation or cause the enzyme to bind to single nicked DNA in spite of post-translational down-regulation is unknown, but recent convincing data demonstrates that the Topo I mediated DNA
10 lesion modification process does occur. The present analogs appear to bind directly to the enzyme before it forms a complex with DNA. Therefore, these compounds inhibit Topo I directly in a very different mechanism than camptothecin derivatives, which only bind a Topo I-DNA intermediate. The overall effect of these Topo I
15 modulators in combination with ionization radiation and/or chemotherapy is that the enzyme is able to bind to damaged DNA and create double-stranded irreversible DNA breaks, induced apoptosis, and kill log-phase as well as confluence arrested human tumor cells which have elevated Topo I levels (i.e., prostate, colon, breast, melanoma and lung cancers).


20 Only a few agents are known to modulate Topo I. The only known drugs that specifically inhibit Topo I are β -lapachone and its derivatives, camptothecin, Hoescht dyes, (which are weak inhibitors) and their respective analogs. Many have attempted to produce various camptothecin and β -lapachone analogs, since the parent drugs have strong toxic side effects at concentrations required for their use alone as
25 chemotherapeutic agents. However, the present invention overcomes the strong cytotoxic effects of these parent compounds, is stabilized in air, water or human serum while still causing sufficient lethality alone or in combination with radiotherapy or chemotherapy against human neoplastic cells.

Summary of the Invention

This invention relates to β -lapachone analogs, substituted at the biologically relevant C-3 position, and their use either alone or to augment radio- and chemotherapy to induce programmed neoplastic cell death.

5 These analogs work using two different mechanisms. First, these analogs can work synergistically with chemo- or radiotherapeutic agents augmenting their potential. Secondly, these analogs work via another mechanism as chemotherapeutic agents.

10 The β -lapachone analogs of the present invention are those having the following structure:



15 where R is $(CH_2)_n-R_1$ where n is an integer from 0-10 and R_1 is hydrogen, an alkyl, an aryl, a heteroaromatic, a heterocyclic, an aliphatic, an alkoxy, a hydroxy, an amine, a thiol, an amide, or a halogen.

20 Specifically, this invention relates to 3-substituted- β -lapachone derivatives particularly 3-hydroxy β -lapachone, 3-alkoxy- β -lapachones, 3-alkyl- β -lapachones and 3-halo- β -lapachones, and their synthesis as well as their use either alone as a chemotherapeutic agent or as an enhancer of the cytotoxic effects of DNA damaging
25 agents (i.e. chemotherapy and radiotherapy). When these analogs are combined with radiation or some other DNA damaging agent which induces DNA strand cisions or cause alkylation damage, a double stranded irreversible break is achieved which results in lethality. These analogs of 3-lapachone require only about one tenth the dose necessary compared to lapachones. 3-Allyl- β -lapachone gives a drug enhanced
30 radiation value (DER value) of 3.6, which is better than any camptothecin derivative (e.g. topotecan) which was administered at 10 times the dose of the 3-allyl- β -lapachone but resulted in a DER value of only 2.7. Therefore, the

surrounding unirradiated tissues would not be affected nor would secondary tumors be induced (they may actually be prevented) in this unirradiated area when using the 3-allyl- β -lapachone.

The administration of efficacious doses of 3-allyl- β -lapachone to human lung, breast, melanoma, prostate, leukemia and colon cancer cells increases the lethal effects of radiation without causing toxicity to surrounding unirradiated normal cells. Tumor selectivity is afforded by elevations of Topo I in cancerous lung, colon, breast, melanoma, prostate and leukemia cells.

Not only does this invention identify and synthesize novel β -lapachone analogs and their anti-neoplastic use, but it discloses a new effective chemical synthesis of these compounds. This novel synthesis is superior over the prior art due to the fact that it produces analogs which are free of impurities and the process is reproducible.

Finally, another use of these 3-substituted β -lapachone derivatives is for the treatment of HIV infection, commonly known as AIDS. Inhibition of Topo I leads to a blockage of the HIV-related reverse transcriptase. The inhibition of reverse transcriptase using β -lapachone has been demonstrated, and it is proposed that the use of the 3-substituted β -lapachones will be for prolonging the survival of AIDS patients and treating their viral infections to prevent the spread of infected cells in the body via the Topo I-mediated inhibition of the HIV-related reverse transcriptase.

These analogs overcome the disadvantages of the prior art (i.e., β -lapachone and camptothecin) due to the fact that these β -lapachone analogs are not cytotoxic to unirradiated, healthy cells surrounding the neoplastic tumor site and require much lower dose levels for effectiveness.

Another improvement over the prior art is that several of these analogs are water soluble derivatives unlike β -lapachone itself and camptothecin. Thus, these analogs are suitable for medical administration and pharmaceutical products.

In addition, another aspect of this invention is the use of 3-substituted β -lapachones to induce programmed cell death. These 3-substituted derivatives induce programmed cell death *without* the use of radiation or additional chemotherapy in cells having elevated Topoisomerase I levels such as cancerous lung, colon, breast, melanoma, prostate, and leukemia cells. Specifically, these cells induce cell death,

inter alia, through apoptotic mechanisms. Thus, these analogs can be used as chemotherapeutic agents themselves.

Brief Description of the Drawings

5 The drawings illustrate the best mode presently contemplated of carrying out the invention.

 In the drawings:

 Fig. 1 is a 0.7% agarose gel showing the inhibition of Topo I by various Topo I modulators;

10 Fig. 1(a) is a 1.0% agarose gel illustrating the initiation of apoptosis by 3-substituted- β -lapachones;

 Fig. 2 is a graph showing percent survival of unirradiated U1-Mel cells as compared to X-irradiated U1-Mel cells with varying levels of 3-allyl- β -lapachone concentrations;

15 Fig. 3 is a graph illustrating percent survival of U1-Mel cells with varying doses of X-irradiation (in cGy);

 Fig. 4 is chart showing percent survival of X-irradiated U1-Mel cells with varying exposure times of 3-allyl- β -lapachone;

 Fig. 5 is a graph illustrating the percent survival of unirradiated compared to
20 X-irradiated U1-Mel cells after the Topo I active drug has been removed, demonstrating irreversible lethality;

 Fig. 6 illustrates the optimal exposure time of a single dose of 3-allyl- β -lapachone by showing the percent survival of unirradiated compared to X-irradiated U1-Mel cells.

25

Detailed Description of the Preferred Embodiment

I. Evidence of Down-Regulation of Topo I and 3-Substituted β -Lapachone I-Induced Topo I Mediated Suicide Pathway Leading to Lethality

5 A. Down-Regulation of Topo I Activity In Normal and Neoplastic Chinese Hamster Embryo Fibroblasts.

To observe changes in Topo I levels with or without ionizing radiation as cells progress toward a neoplastic state, Topo I activities were measured in unirradiated or X-irradiated confluence-arrested normal diploid CHEF/18 cells and their neoplastic derivatives: 21-2, 21-2M3, and T30-4. CHEF cells were treated with 400 cGy, an equitoxic dose resulting in 20% survival for each cell type. Topo I activities of unirradiated and X-irradiated CHEF cells at 4 h posttreatment were assayed in 4 separate experiments. Topo I enzymatic and survival recovery data following radiation for all four cell types are summarized in Table 1.

15

TABLE 1

Down-Regulation of Topo I Activity in Normal
and Neoplastic Chinese Hamster Embryo
Fibroblast (CHEF) Cells.¹

20

Cell Type (Malignancy)	X-Ray Dose (cGy)	Topo I ¹ Enzyme Activity	Extent of Topo I ¹ Down-Regulation	%Survival ³ Enhancement
CHEF/18	0	4.0 \pm 0.5	2X	22 \pm 2
(0/42)*	400	2.0 \pm 0.1		
21-1	0	4.1 \pm 1.0	3 X	25 \pm 4
(0/42)*	400	1.2 \pm 0.04		
T30-4	0	11.4 \pm 0.1	29X	60 \pm 7
(42/42)*	400	4.0 \pm 0.01		
21-2M3	0	6.7 \pm 0.1	10X	45 \pm 12
(18/42)*	400	0.7 \pm 0.02		

30

¹Topo I enzyme activity, defined as the loss of 60% supercoiled DNA substrate (i.e., 300 μ g p36B4 plasmid) per mg protein per minute, for X-irradiated

and unirradiated cells at 4 h post-irradiation was determined from the slope of the linear portion of the Topo I activity curves in Figures 2A through 2D (see Appendix. 13). *Indicates number of tumors formed in nude mice (43). ²Extent of down-regulation of Topo I enzyme activity following 400 cGy was calculated by the ratio of Topo I enzyme activities of unirradiated to X-irradiated cells. ³Survival enhancement values for CHEF cells were previously determined using colony forming assays (33.42.43) by subtracting the survival of cells immediately following ionizing radiation (i.e., no posttreatment) from the survival of irradiated cells after 8 h recovery.

10

A gradual increase in Topo I activity was noted as cells progressed to neoplasia. When unirradiated CHEF/18 cells were compared to its neoplastic derivatives, higher enzyme activities in the order of: CHEF/18 < 21-2 < 212M3 < T30-4 were found. Unirradiated T30-4 cells, the most malignant cell type, contained nearly 3-fold higher Topo I levels than unirradiated CHEF/18 cells. Following radiation, Topo I levels were dramatically down-regulated and the extent of down-regulation increased as cells became more neoplastic. Topo I down-regulation correlated with the capacity of damaged cells to recover from radiation; cells which recovered well from radiation damage also demonstrated a greater capacity to down-regulate Topo I (see Table 1).

20

Camptothecin, a specific Topo I inhibitor, was used to prove that Topo I and not Topo II activities were assayed. Extracts from unirradiated CHEF/18 cells and its neoplastic derivatives (i.e., 21-2M3 and T30-4) containing Topo I activities were inhibited 100% by camptothecin (50 nM-1 μ M) at 30 minutes and 2 hours. Administration of lower levels of camptothecin (between 1 nM and 10 nM) resulted in partial inhibition of Topo I. Enzymatic activity was observed in controls and after administration of 500 nM m-AMSA, a potent Topo II inhibitor. Topo II levels were not detected in confluence-arrested human or CHEF/18 cells. This is consistent with previous results showing that Topo II is not present in quiescent cells.

30

B. Temporal Regulation of Topo I in X-irradiated CHEF Cells.

Topo I activity was investigated in CHEF cells after various times of recovery following radiation. Normal CHEF/18 and transformed 21-2M3 cells were unirradiated or X-irradiated (400 cGy) and incubated at 37°C for various times.

- 5 Topo I activity decreased in both CHEF/18 and 21-2M3 cells within 5 minutes and remained low for the next 12 hours. Topo I activities did not recover for nearly 18 hours after radiation, at which time only 75% control levels were noted. Topo I levels recovered completely in 24 hours.

10 *C. Down-Regulation of Topo I Activities In Human Cells After Ionizing Radiation.*

- Topo I activity in human malignant melanoma (U1-Mel) cells was examined before and after X-irradiation. As in CHEF cells, Topo I activity decreased in X-irradiated U1-Mel cells within 5 minutes post-irradiation. The lowest Topo I level
15 was noted at 45 minutes post-irradiation. Topo I activity in irradiated U1-Mel cells remained low until 65 minutes, when enzyme activity increased to roughly 80% unirradiated control levels. Unlike CHEF cells which took 24 hours for Topo I levels to recover, Topo I activity within X-irradiated U1-Mel cells returned to levels comparable to control cells within 65-120 minutes. We also examined Topo I activity
20 in U1-Mel cells following various doses of ionizing radiation at 45 minutes. Topo I activity was gradually down-regulated by increasing doses of radiation up to 450 cGy. At doses of radiation greater than 450 cGy, Topo I activity remained at 1-2% control levels.

- Topo I down-regulation was observed in a variety of human normal and tumor
25 cells following equitoxic doses of radiation giving 10% survival, with similar temporal down-regulation as noted in U1-Mel cells (see Table 2). Higher Topo I activities were noted in unirradiated human tumor (U1-Mel, HTB-152, and HEp-2) compared to human normal (GM2936B, GM2937A, and IMR-90) cells. Topo I activities in X-irradiated human tumor cells were, however, down-regulated to a
30 greater extent than in human normal cells. In fact, Topo I levels following radiation decreased in X-irradiated human tumor cells to far lower levels than observed in irradiated normal cells ($p < 0.1$, Student's *t* test). As in CHEF cells, the capacity to

down-regulate Topo I after radiation correlated with enhanced survival in various human cells after radiation.

It is logical that if down-regulation of Topo I occurs in cells damaged by irradiation in order to repair incised DNA breaks, then cells whose DNA is damaged by chemotherapy will also down-regulate Topo I in the same manner; for example, it has been demonstrated that UV-irradiation also down-regulates Topo I. Furthermore, it has been demonstrated that presubstituted β -lapachone compounds will increase lethality alone (via apoptosis) and in combination with radiotherapy or chemotherapy by preventing Topo I down-regulation and causing the Topo I mediated suicide pathway discussed above and illustrated in Pathway 1.

TABLE 2
Decreased Topo I Activity Following Ionizing Radiation in Human Cells

Cell Type	X-Ray Dose ¹ (cGy)	Topo I ² Enzyme Activity	Extent of Topo I ¹ Down-Regulation	%Survival ³ Enhancement
<u>Tumor Cells</u>				
U1-Mel	0	10.8 \pm 0.4	23X	45 \pm 3.4
	450	0.5 \pm 0.05		
HTB-152	0	9.0 \pm 0.2	23X	25 \pm 0.7
	400	0.4 \pm 0.1		
HEp-2	0	7.1 \pm 0.02	19X	37 \pm 1.5
	400	0.4 \pm 0.3		
<u>Normal Cells</u>				
GM29236B	0	2.5 \pm 0.2	2X	17 \pm 1.3
	290	0.8 \pm 0.01		
GM2937A	0	2.5 \pm 0.3	3X	15 \pm 3.2
	300	0.9 \pm 0.02		
IMR-90	0	4.2 \pm 0.5	4X	21 \pm 2.8
	350	1.0 \pm 0.01		

¹Topo I enzyme activities, defined as the loss of 60% supercoiled DNA substrate (i.e., 300 µg p36B4 plasmid) per mg protein per minute, for X-irradiated and unirradiated cells at 45 min post-irradiation were determined from the slope of the linear portion of the Topo I activity curves. Experiments were repeated three
5 times in duplicate.

²Extent of down-regulation of Topo I enzyme activity following equitoxic doses of ionizing radiation was calculated by the ratio of Topo I enzyme activities of unirradiated to X-irradiated cells.

³Survival enhancement values were determined by performing colony forming
10 assays and values were calculated by subtracting the survival of cells immediately following ionizing radiation (i.e., no posttreatment) from the survival of irradiated cells after 4 h recovery (33,42).

D. Regulation of Topo I mRNA and Protein Levels After Radiation.

15 Northern blot analyses using ³²P-labeled human Topo I cDNA revealed that Topo I mRNA levels (MW: 4.2 Kb) were unchanged in CHEF and U1-Mel cells before and after ionizing radiation. Blots were also probed for 36B4 expression and a Northern loading control was unaltered with radiation as previously observed. Since Topo I mRNA levels were not altered by radiation, we then examined changes
20 in Topo I protein levels in CHEF and U1-Mel cells. U1-Mel and CHEF cells were treated with or without ionizing radiation, nuclear protein extracts were then analyzed on 7% SDS-PAGE gels, and Topo I protein levels were detected by Western blot analyses as previously described. Topo I protein levels in CHEF cells or in U1-Mel cells were not altered by ionizing radiation. Topo I protein levels also remained
25 unchanged in U1-Mel cells at various times post-irradiation (450 cGy) as monitored by Western blot analyses and Topo I immunoprecipitation.

E. Topo I Down-Regulation Occurs Via ADP-Ribosylation.

The mechanism by which Topo I activity was down-regulated by radiation was
30 then explored using inhibitors of poly (ADP-ribose) transferase (ADPRT), an enzyme which adds poly (ADP-ribose) units to proteins. Topo I activity is modulated by phosphorylation, which activates the enzyme at the G₁/S border, and by poly

(ADP-ribosylation) which inactivates the enzyme. The effects of 3-ABA and PD 128763 (hereafter called PD 1), a more potent inhibitor of ADPRT, were examined on Topo I activity following radiation. The administration of 5 μ M 3-ABA or 0.5 μ M PD 1 to U1-Mel cells 1 hour prior to, or for 4 hours immediately after, 400 cGy prevented the down-regulation of Topo I enzyme activity. A five hour exposure of unirradiated U1-Mel cells to 0.5 μ M PD1 caused only a slight loss of viability ($92 \pm 1.6\%$ viability) of unirradiated U1-Mel cells, but decreased the survival of X-irradiated U1-Mel cells to $5 \pm 0.2\%$. The administration of 0.5 μ M PD 1 to U1-Mel cells restored Topo I activity equivalent to that observed in unirradiated U1-Mel cells. The fact that Topo I activities were down-regulated following UV-irradiation (10 J/M^2) was noted, and the administration of 0.5 μ M PD 1 also prevented this down-regulation.

Since no apparent shift in Topo I protein molecular weight was noted in the analyses above, changes in Topo I protein were monitored using two-dimensional gel electrophoresis and Western blot analyses in unirradiated and X-irradiated (450 cGy) U1-Mel cells. Slight shifts in both molecular weight (approximately 2-5 kDa) and isoelectric point (a shift from pI: 8.2 to pI: 8.34) of Topo I from X-irradiated U1-Mel cells were noted using several known protein spots (e.g., β -actin, alpha-tubulin), as references. These data suggest that if poly (ADP-ribosylation) by ADPRT of Topo I is responsible for its down-regulated activity, the substitution must be a mono-ADP-ribosylation, possibly at a site within Topo I which is normally phosphorylated.


Based on the aforementioned data, it appears that the 3-substituted β -lapachones interact with Topo I after exposure to irradiation, chemotherapy or any DNA damaging agent and is able to interfere with Topo I down-regulation thereby increasing DNA damaging agent lethality.

II. Novel Synthesis of 3-Substituted β -Lapachone Derivatives.

Lapachol (1) is a natural product known since 1858 and is the most abundant quinone found in the heartwood of several genera Bignonoceae. Its name is derived from the "lapacho" tree (*Tabebuia avellanedae* Lr. ex. Griseb), a tree found mainly in subtropical and tropical South America (Northern Argentina, Paraguay, and

Brazil). The yellow lapachol confers its color to the wood where it is sometimes visible in yellow deposits. The structure of lapachol has been known and its antitumor properties raised considerable interest in the past, having reached the stage of clinical trials. When treated with sulfuric acid it cyclizes to β -lapachone (2a), also present in the heartwood of the lapacho tree. Lapachol can also be converted into 3-bromolapachone (2b), and the latter can be converted in a two-step sequence into 3-hydroxy- β -lapachone (2c). The alcohol group of 3-hydroxy- β -lapachone is amenable to etherification when treated with alkyl halides in dry dimethyl sulfoxide in the presence of potassium hydroxide powder. It was found that it is possible to alkylate the 3-hydroxy residue of 3-hydroxy- β -lapachone without ring opening of the β -lapachone structure by avoiding alkaline reaction media which are usually necessary in etherification or esterification procedures. It was thus possible to obtain methyl, benzyl, and allyl ethers (2d) of 3-hydroxy- β -lapachone, as well as β -ethoxycarbonyl derivatives. (See Synthesis 1). In synthesis 1 and 2 R is as previously defined herein.

SYNTHESIS 1



What makes this approach attractive is that lapachol is a relatively abundant natural product. In the heartwood of the abundant South American lapacho tree, its content varies between 3% (in subtropical South America) to 7% (in tropical South America). These values should be compared with those for camptothecin (0.01% in

the stem wood of *C. accuminata*) or in taxol (0.02% in the bark of the American western yew).

When the silver salt of lapachol is alkylated (see Synthesis 2) the main products are the O-alkyl ethers of lapachol (3).


5

SYNTHESIS 2

10

15

20



However, a very small amount of 3-substituted β -lapachones (4), obtained in 4% yield, are also formed. This side reaction takes place with the silver salt but not with the potassium or sodium salts of lapachol. The 3-substituted β -lapachones can be separated from the main reaction product 3 by extraction with a 5% bisulfite solution.

Purification of 3-substituted β -lapachones requires a final chromatographic step on silica gel, using variable amounts of methanol in methylene chloride. The isolated amounts of 3-substituted- β lapachones are now ready for biological assays on Topo I enzyme assays and cell culture studies. The precise details of this general procedure are outlined below.

A. General procedures of the Synthesis of 3-O-alkyl- β -lapachone ethers

3-Hydroxy- β -lapachone- (258 mg, 1 mmol) was added to a mixture of dry and finely powdered potassium hydroxide (224 mg, 4 mmol) in 2 ml of dimethyl sulfoxide. The mixture was stirred for 10 min at 20°C, when the alkyl halide (2 mmol) was added. Stirring was continued during 30 min, water (20 ml) was then added, and the aqueous solution was then extracted with methylene chloride (3 x 15 ml). The pooled extracts were washed with water (3 x 30 ml), dried (Na_2SO_4) and evaporated to dryness. The residue was dissolved in a small volume of methylene chloride, adsorbed on a silica gel column prewashed with the same solvent, and the orange band was eluted with 5% methanol in methylene chloride. The eluates were evaporated to dryness, and the crystalline residue was recrystallized from ethanol.

3-Methoxy- β -lapachone (2a) was obtained in 60% yield using methyl iodide; mp $215 \pm 216^\circ\text{C}$; $^1\text{HNMR}$ (CDCl_3 , ppm) 7.9 (dd, 2H), 7.6 (dd, 2H), a,b arom H; 3.4 (s, 3H) OCH_3 ; 2.7-2.2 (dd, 2H) CH_2 -4; 1.90 (m, H) H-3; 1.6 (s, 3H), 1.3 (s, 3H) CH_3 -2.

3-Benzoyloxy- β -lapachone (2b) was obtained in 6% yield using benzyl bromide; mp $198 \pm 199^\circ\text{C}$; $^1\text{HNMR}$ (CDCl_3 , ppm) 7.9 (dd) 7.6 (dd); arom H; 7.3 (s, 5H) Bn; 4.6 (s, 2H) CH_2 Bn; 2.7 (dd) 2.15 (ddd) CH_2 -4; 1.90 (m, H) H-3; 1.6 (s), 1.3 (s) CH_3 -2.

3-Ethoxycarbonylmethyloxy- β -lapachone (2c) was obtained in 23% yield using ethyl bromoacetate; mp $156 \pm 157^\circ\text{C}$; $^1\text{HNMR}$ (CDCl_3 , ppm) 7.6 (dd)-7.6 (dd) arom H; 4.25 (q, 2H) CH_2CH_3 ; 4.10 (s, 2H) CH_2CO ; 2.7-2.15 (dd) CH_2 -4; 1.95 (m, H) H-3; 1.6 (s)-1.3(s) CH_3 -2.

3-Allyloxy- β -lapachone (2d) was obtained in 55% yield using allyl bromide; mp $125 \pm 126^\circ\text{C}$; $^1\text{HNMR}$ (CDCl_3 , ppm) 7.9-7.6 (dd) arom H; 5.88 (m, H) $\text{CH}=\text{}$; 5.30 (m, 2H) $\text{CH}_2=\text{}$; 4.0 (m, 2H) CH_2O ; 2.77-2.15 (dd) CH_2 -4; 2.0 (m) H-3; 1.6 (s)-1.3 (s) CH_3 -2.

B. General Procedure of Synthesis of 3-alkyl- β -lapachone derivatives

Lapachol (1g) is dissolved in 200 ml of a 0.01 N ammonium hydroxide solution with stirring and warming at 40°C. After the solid went into solution, the latter is filtered to eliminate impurities, and the excess ammonia is chased away with

a stream of nitrogen so as to bring the solution to pH ca. 7.0. An aqueous solution of silver nitrate is then slowly added until the intense red color of the solution is discharged. The red precipitate is filtered, washed with distilled water, then with 95% ethanol, and finally with ether. The red silver salt (1.4 g) is dried in vacuo and
5 grinded in a porcelain mortar. It is then suspended in 100 ml of dry ether, and a ten-fold amount of the alkyl halide is added. The mixture is stirred and heated under reflux until the red silver salt disappeared and a dark brown precipitate of silver halide was formed (reflux times varied between 0.5 hr-12 hr, depending on the halide). Conversely, the reaction can be cooled instead of heated. The silver halide
10 was filtered and washed with ether (20 ml). The filtrates were pooled, washed with 1M sodium hydroxide (3 x 20 ml) to eliminate residual lapachol, then with water (5 x 10 ml) to neutral, dried (Na_2SO_4), and evaporated to dryness. The oily residue was dissolved in a small volume of chloroform, and adsorbed on a silica gel column prewashed with the same solvent. The column was washed with this eluant which
15 eluted the O-alkyl ethers (3) as yellow bands, followed then by methylene chloride. Derivatives 3-alkyl- β -lapachol (4b) and 3-methyl- β -lapachol (4a) were eluted with the latter as orange bands; 3-ethoxy carbonyl methyl- β -lapachol (4c) was eluted with 1% methanol in methylene chloride; and 3-(2'-Hydroxyethyl)- β -lapachone (4d)) and 3'-(2'-Aminoethyl)- β -lapachone (4e) with 5% methanol in methylene chloride. The
20 eluate containing the orange band was evaporated to dryness, and the residue crystallized from the corresponding solvent.

3-Methyl- β -lapachone (4a) was obtained in 1% yield using methyl iodide; mp $179 \pm 180^\circ\text{C}$ (from cyclohexane); $^1\text{HNMR}$ (CDCl_3 , ppm) 7.9-7.6 (dd) arom H; 2.7-2.15 (dd) CH_2 -4; 1.9 (m) H-3; 1.6 (s)-1.3 (s) CH_3 -2; 1.07 (d) CH_3 -3.

25 3-Allyl- β -lapachone (4b) was obtained in 4% yield using allyl bromide; mp $147 \pm 148^\circ\text{C}$ (from hexane); $^1\text{HNMR}$ (CDCl_3 , ppm) 8.0-7.6 (dd) arom H; 5.15 (m, H), 5.85 (m, 2H) vinyl; 2.8, 2.15 (dd) CH_2 -4; 2.45 (m, CH_2 allyl); 1.85 (m, H) H-3; 1.6 (s, 3H), 1.3 (s, 3H) CH_3 -2.

3-Ethoxy carbonyl methyl- β -lapachone (4c) was obtained in 1% yield using
30 ethyl bromoacetate; mp $118 \pm 119^\circ$ (from hexane); $^1\text{HNMR}$ (CDCl_3 , ppm) 7.9-7.6 (dd), arom H; 2.8-2.15 (dd) CH_2 -4; 4.25 (q) CH_2CH_3 ; 3.55 (s) CH_2CO ; 1.9 (m) H-3; 1.61-1.4 (s) CH_3 -2; 1.3 (q) CH_3CH_2 .


3-(2'-Hydroxyethyl)- β -lapachone (4d) was obtained in 2% yield using 2-bromoethanol; mp $205 \pm -206^\circ$ (from ethanol-water); ^1H NMR (CDCl_3 , ppm) 7.9-7.6 (dd) arom H; 3.8 (m, 2H) CH_2OH ; 3.6 (b, H) OH; 2.8-2.15 (dd) CH_2-4 ; 2.0 (m) H-3; 1.70 (m), 1.60 (s), 1.40 (s) (8H) CH_3-2 , CH_2-1' .

- 5 3-(2'-Aminoethyl)- β -lapachone (4e) was obtained in 1% yield using 2-bromoethylamine; mp $216 \pm -217^\circ\text{C}$ (from ethanol water); ^1H NMR (CDCl_3 , ppm) 7.9-7.6 (dd) arom H; 5.0 (b, 2H) NH_2 , 3.02 (m, 2H) CH_2NH_2 ; 2.8-2.15 (dd) CH_2-4 ; 2.0 (m) CH_2-1' ; 1.85 (m, H_3); 1.6-1.3, CH_3-2 .

This new protocol and purification synthesis for the 3-substituted compounds leads to functional yields and relative 100% purity. This synthesis is also reproducible. Therefore, this enables the use of these 3-substituted compounds as radio-and/or chemotherapy augmentors.

As used herein the term alkyl represents a straight chain or branched hydrocarbon from 1-10 carbons and all its isomeric forms such as methyl, ethyl, propyl, isopropyl, butyl, isobutyl, pentyl, etc. The aliphatic can be a saturated, unsaturated or polyunsaturated chain. Representative aliphatic groups are the paraffin, olefin and acetylene hydrocarbons and their derivatives.

The term aryl denotes an organic radical derived from an aromatic hydrocarbon by the removal of a hydrogen atom. Aryl signifies a phenyl, or an alkyl, nitro, or halo-substituted phenyl group. Specifically, the term "substituted" as applied to "phenyl" refers to phenyl which is substituted with one or more of the following groups: alkyl, halogen (that is fluorine, chlorine, bromine or iodine), nitro, cyano, trifluoromethyl and the like. The phenyl may be substituted with any electron donating or withdrawing groups. The term alkoxy signifies the group O-alkyl. The hydroxy group may exist as -OH or be esterified with a lipid group and have water soluble functionality. The amine may exist as a primary, secondary or tertiary amine and its derivatives. The amine may exist as $-\text{NH}_2$, $-\text{NHX}$ where X is an alkyl from 1-10 carbons in all isometric forms, and



The amide may have lipid or water soluble functionality and may have an additional CH₂ before the hydroxy. The amide can be a primary or secondary amide in the form -CONH₂ or -CONHX where X is an alkyl. The amide may also be in the form of a primary or secondary amide in the form -CH₂CH₂CONH₂ or -CH₂CH₂CONHX where X is an alkyl. The term thiol signifies -SH or -SX where X is an alkyl from 1-10 carbon atoms in all isometric forms. The halogen may be chlorine bromine, fluorine or iodine. A heteromatic is a substitute aromatic compound examples being pyridine, pyrrole, pyrimidine, imidazole, oxazole, thiophene, thiazole, morpholine, pyrezine, triazine and the like. Heterocyclics are substituted cyclic compounds.

10

III. EFFECTIVENESS OF 3-SUBSTITUTED β -LAPACHONE ANALOGUES

A. Topo I Enzyme Inhibition Measurements and Induction of Apoptosis In HL60 Cells Following 3-substituted β -Lapachone Analogs.

After the synthesis of 3-substituted β -lapachone derivatives (noted above in part II), the ability of 3-substituted β -lapachone derivatives to: (a) modulate purified human Topo I; and (b) cause apoptosis using HL60 (human promyelocytic leukemia) cells, in which diagnostic 180 kbp DNA laddering and loss of intact genomic DNA are observable indicators of apoptosis.

It was demonstrated that all 3-substituted compounds, including the water-soluble 3-hydroxy- β -lapachone, inhibited Topo I in a strict, sequence-specific manner, similar to that of β -lapachone. This was shown using the previously described DNA unwinding assays found in Boothman et.al., "Inhibition of Potentially Lethal DNA Damage Repair in Human Tumor Cells by β -lapachone, an Activator of Topo I" Cancer Res. 49:605-612, (1989). Figure 1 demonstrates the inhibitory effects of 3-allyl- β -lapachone on the activity of human Topo I when the enzyme and drug were incubated for 5 minutes at 37°C prior to the addition of supercoiled pBR322 DNA substrate. The compounds were not effective Topo I inhibitors if the drug and DNA were incubated first, and the DNA unwinding reactions initiated by the addition of Topo I. The reverse was true for camptothecin derivatives. Topo I was inhibited by as little as 10 nM 3-allyl- β -lapachone and an estimated K_i of approximately 100 nM was determined. Complete inhibition of Topo I was observed

at 1 μ M for all 3-substituted β -lapachone compounds, including the water-soluble, 3-hydroxy- β -lapachone derivative.

For comparison, an estimated K_i of 350 nM for camptothecin to Topo I was measured under reaction conditions in which the drug was added to the DNA before the enzyme. The Topo I inhibitory effect of 3-allyl- β -lapachone was greater than
5 β -lapachone; a K_i of 150 nM was calculated for β -lapachone.

It was discovered that β -lapachone is highly instable in air, even more so in water, and breaks down over time. There is no observance of a similar break-down reaction with the 3-substituted β -lapachone compounds in air. In addition, 3-allyl- β -
10 lapachone was more stable in water than β -lapachone, but both compounds lose stability over time when solubilized in water or DMSO (i.e., >4 days with 3-allyl- β -lapachone; <2 days with β -lapachone). In conclusion, the results with the 3-substituted β -lapachone compounds indicate that they are very effective Topo I inhibitors which have activities greater than that of camptothecin or its analogs.

15 The effect of 3-substituted β -lapachone derivatives on the initiation of apoptosis using HL60 (human promyeloid leukemia) cells was then observed, where 180 kbp DNA laddering and loss of intact genomic DNA are diagnostic for the induction of apoptosis. The methodology was illustrated in Bertrand et.al., "Apoptosis and its Modulation in Human Promyelocytic HL-60 cells treated with DNA
20 Topoisomerase I and II Inhibitors", Blood 81: 1359-1368, 1993. For all 3-substituted β -lapachone derivatives, apoptosis in HL60 cells has been observed (see Figure 1(a) for apoptosis with 3-hydroxy- and 3-allyl- β -lapachone) after 4 hour treatment with 0.5-1.0 μ M; complete apoptotic responses were observed at 1.0 μ M and apoptosis was observed after as little as one hour after treatment. Only minor
25 differences in effective apoptotic-inducing concentrations between 3-substituted compounds were observed. The induction of apoptosis and inhibition of Topo I activity after various concentrations of 3-substituted β -lapachone compounds compared to that occurring after β -lapachone or camptothecin exposures was then compared. For all tests of 3-substituted- β -lapachone compounds, greater apoptosis
30 and Topo I inhibition (at lower efficacious doses) were found compared to the β -lapachone prepared by Ciba Geigy. For each compound, the induction of apoptosis correlated well with effective Topo I inhibitory dose-responses, with the 3-substituted

compounds being more effective at inhibiting Topo I and initiating apoptosis at 20-40% lower concentrations than β -lapachone. In contrast, α -lapachone or lapachol (compounds also dissolved in DMSO and used at various concentrations) did not inhibit Topo I as previously described, nor did they induce apoptosis.

5

B. Radiosensitization Effects of 3-Allyl- β -lapachone.

The potential radiosensitizing effects of the more stable β -lapachone derivative, 3-allyl- β -lapachone, against human radioresistant melanoma (U1-Mel) cells was then analyzed (Figures 2-6):

10

(i.) Optimal Concentration of 3-Allyl- β -lapachone.

The toxic effects of 3-allyl- β -lapachone on confluence-arrested human U1-Mel cells were studied. Concentrations of 3-allyl- β -lapachone up to 25 μ M were less toxic to unirradiated U1-Mel cells when treated for 4 hours than similar treatments with β -lapachone. An optimal concentration range of 4-40 μ M 3-allyl- β -lapachone was determined from the data in Figure 2, in which relatively little toxicity to unirradiated U1-Mel cells were noted. Therefore, we chose 10 μ M β -lapachone as an optimal dose for continued studies. In fact, in other experiments a very large range of concentrations of 3-allyl- β -lapachone (10-40 μ M) with 400 cGy appeared to be optimal, causing minimal lethal effects to unirradiated confluence-arrested U1-Mel cells, while greatly enhancing the lethality of X-irradiated U1-Mel cells.

15

20

(ii.) X-ray Dose-Response Following 3-Allyl- β -lapachone.

Using an optimal concentration of 3-allyl- β -lapachone (10 μ M), its ability to enhance the lethal effects of various doses of ionizing radiation was investigated (Figure 3). Irradiated U1-Mel cells which were allowed to repair for 4 hours in the absence of drug increased survival as compared to cells which were not allowed to repair (i.e., cells given a no posttreatment, NPT). Addition of 3-allyl- β -lapachone immediately after radiation resulted in synergistic enhancement of lethality. Enhanced lethality of 3-allyl- β -lapachone was directly proportional to the amount of damage caused by radiation; the higher the dose of radiation the greater the enhancement (Figure 3). 3-Allyl- β -lapachone resulted in greater dose enhancement ratios (DERs)

25

30

(3.6 ± 0.3) compared to optimal doses of β -lapachone (at $4\mu\text{M}$, DER: 1.4 ± 0.4) or camptothecin (at $4\mu\text{M}$, DER: 2.1 ± 0.6) at 10% survival levels (see Table 3).

(iii.) Timing of 3-Allyl- β -lapachone Exposure for Radisensitizing Effects.

5 Using an optimal dose of 3-allyl- β -lapachone ($10\mu\text{M}$) and an optimal drug exposure time of 4 hours, the effect of variations in timing of 3-allyl- β -lapachone addition was examined (Figure 4). When confluence-arrested U1-Mel cells were exposed to $10\mu\text{M}$ 3-allyl- β -lapachone for 4 hours prior to radiation, no affect on survival recovery was noted. In contrast, U1-Mel cells treated with $10\mu\text{M}$
10 3-allyl- β -lapachone during or immediately after radiation resulted in synergistic enhancement of lethality. Treatment of unirradiated cells with $10\mu\text{M}$ 3-allyl- β -lapachone did not significantly effect survival. Similar timing results with other Topo I-active drugs (i.e., camptothecin, β -lapachone, topotecan and 9-aminocamptothecin) have been described. Thus, these data are consistent with other
15 radiosensitizing effects observed by Topo I inhibitors.

(iv.) Reversibility of 3-Allyl- β -lapachone Radiosensitization.

Confluence-arrested U1-Mel cells were unirradiated or X-irradiated (400 cGy) and exposed to 0.2% FCS-DME, $10\mu\text{M}$ 3-allyl- β -lapachone for 4 hours. After
20 incubation in the Topo I active drug, cells were washed and then incubated for an additional time in DME + 10% FCS without 3-allyl- β -lapachone (Figure 5) to test for reversibility of lethality. Another set of confluence-arrested U1-Mel cells were irradiated and plated immediately for survival to prevent repair (i.e., NPT). Survival levels were corrected for loss of plating efficiency due to prolonged confluence-arrest
25 and drug exposures alone. Irradiated cells not exposed to 3-allyl- β -lapachone, but allowed to recover, increased their survival from 50% to 80% in 8 hours. X-irradiated cells exposed for 4 hours with $10\mu\text{M}$ 3-allyl- β -lapachone did not repair after drug removal. The irreversible effects of post-irradiation exposure to $10\mu\text{M}$ 3-allyl- β -lapachone were similar to that those observed with β -lapachone,
30 camptothecin, topotecan or 9-aminocamptothecin.

(v.) Kinetics of Radiosensitization by 3-Allyl- β -lapachone.

The optimal exposure time of a single dose of 3-allyl- β -lapachone (10 μ M) that could enhance the efficacy of radiation was then determined (Figure 6). U1-Mel cells were X-irradiated (400 cGy) and allowed to repair in the absence of presence
5 of 10 μ M 3-allyl- β -lapachone. A set of X-irradiated U1-Mel cells were trypsinized and replaced immediately after radiation to determine the survival of cells not allowed to repair (i.e., NPT). Longer exposure times (i.e., 4-10 hours) of unirradiated U1-Mel cells to 3-allyl- β -lapachone did not enhance cytotoxicity. An exposure time of 4-6 hours was optimal.

10

C. Comparison of Topo I-active Drugs to Potentiate Ionizing Radiation Lethality.

Several Topo I modulating drugs were collected (some of which are currently in clinical trials) and used to perform five separate types of experiments (in
15 accordance with the experiments listed in i-v above). For each drug: (a) the optimal drug dosage which caused radiosensitization was determined; (b) determined dose enhancement responses; (c) evaluated optimal timing of drug addition; (d) the influence of drug exposure on unirradiated and X-irradiated cells was investigated; and (e) determined whether or not drug-induced radiosensitization was reversible or
20 irreversible. Topo I inhibitors of interest included β -lapachone, camptothecin, 3-allyl- β -lapachone, 9-aminocamptothecin, and topotecan. The results are summarized in Table 3.

Posttreatment addition of 3-allyl- β -lapachone was the most effective agent at increasing the sensitivity of radioresistant human U1-Mel cells to X-irradiation.

25

TABLE 3
Effect of Topo I Modulators On X-Ray-Sensitivity
Of Human U1-Mel Cells¹.

5	Treatment (4 Hr. Post)	Viability (%) of Unirradiated Cells	D _q 3 (cGy)	D ₀ 3 (cGy)	10%	DER Values ¹ 1%	n
	DME, No Post	100	290±8	210±13	1.0±0.02	1.1±0.02	66
10	DME, 0.2% FCS	100	412±14	342±10	---	---	66
	μM β- lapachone	79±4	152±21	154±8	1.5±0.3	1.4±0.41	36
15	10 μM 3- Allyl-β- lapachone	89±3	56±4	54±2	4.2±0.2	3.6±0.3	16
20	4 μM Camp- tothecin	54±2	132±2	162±7	1.8±0.1	2.1±0.6	22
	<i>Camptothecin Derivatives</i>						
	2 μM Topotecan	95±3	190±6	186±4	1.4±0.6	1.6±0.4	8
25	4 μM Topotecan	92±4	127±5	132±5	1.6±0.2	1.7±0.3	8
	100 μM Topotecan	63±7	100±3	113±2	2.2±0.2	2.7±0.2	10
30	10μM 9- Amino- 20(RS)-	67±5	110±2	109±2	2.5±0.3	2.9±0.4	12
	10μM 10,11-	62±7	345±3	254±5	1.2±0.2	1.4±0.1	8
35	Methylene- dioxy-100μM 9- Glycinamido- 20(RS)-	61±3	352±4	278±6	1.1±0.1	1.4±0.4	8
40	10μM 20-Amino- 20(RS)- Deoxy(RS)	64±3	210±5	197±8	1.4±0.6	1.8±0.3	8

¹ 3-allyl-β-lapachone administered at 10 μM demonstrated greater dose
enhancement ratios (DERs) (3.6 ± 0.3) compared to optimal doses of β-lapachone (at
4 μM, DER: 1.4 ± 0.4) camptothecin (at 4 μM, DER: 2.1 ± 0.6), Topotecan (at

10 μM , DER: $2.3 \pm .02$), and 9-aminocampotecin (at 10 μM , DER: 2.9 ± 0.4)
 These data indicate that 3-substituted β -lapachone derivatives are more effective
 radiosynthetizers than camptothecin derivatives or β -lapachone.

5 IV. Inhibition of HIV-Related Reverse Transcriptase Activity By β -Lapachone

The effects of Topoisomerase I (Topo I) inhibition caused by β -lapachone on
 the activity of HIV-related reverse transcriptase activity. In these reactions, reverse
 transcriptase is measured by the level of incorporation of (^3H) dCTP into cDNA by
 the enzyme using a poly A⁺ mRNA as substrate. Reactions were prepared containing
 10 increasing β -lapachone concentrations, with or without the administration of Topo I.
 cDNA resulting from the reaction products were then analyzed by trichloroacetic acid
 (TCA) precipitation onto glass fiber filters. Radioactive cDNA resulting from the
 reverse transcriptase reaction was then measured via liquid scintillation counting. The
 higher the cpm on the glass fiber filters, the higher the enzyme activity. The results
 15 are demonstrated below:

TABLE 4

	Condition*	cpm On Glass Fiber Filter	
20	1. AMV Control +	57,166	$\pm 9,242$
	2. AMV + 4 μM β -lapachone	61,812	$\pm 6,259$
	3. AMV + 10 μM β -lapachone	67,942	$\pm 15,969$
	4. AMV + 50 μM β -lapachone	44,838	$\pm 2,956$
	5. AMV + 100 μM β -lapachone	59,358	$\pm 8,616$
25	6. AMV + Topoisomerase I (Topo I)	14,375	$\pm 2,342$
	7. AMV + Topo I + 4 μM β -lapachone	14,783	$\pm 6,351$
	8. AMV + Topo I + 10 μM β -lapachone	10,092	$\pm 1,862$
	9. AMV + Topo I + 50 μM β -lapachone	9,431	$\pm 4,458$
	10. AMV + Topo I + 100 μM β -lapachone	7,245	$\pm 1,811$
30	11. No Addition Control (Background)	3,069	$\pm 1,235$


* AMV was used at 1 unit/reaction.

+ DMSO was used at a concentration equivalent to those used in β -lapachone
 treatments.

The results indicate that β -lapachone does not influence the enzymatic activity of AMV reverse transcriptase alone. In contrast, addition of Topo I caused a decrease in reverse transcriptase activity, which is then greatly augmented by the concomitant addition of β -lapachone. This decrease by Topo I alone could be due to the rather high levels of Topo I (3 units/reaction) used in the reaction conditions. The results on the inhibition of reverse transcriptase by β -lapachone, only in the presence of Topo I, are consistent with previous results which demonstrated that β -lapachone inhibits this enzyme and prolongs the life of Rous sarcoma virus infected chickens (Schaffner-Sabba K., Schmidt-Ruppin KH, Wehrli W, Schuerch AR, and Wasley JWF. " β -lapachone: Synthesis of Derivatives and Activities in Tumor Models.," J. Med. Chem 27: 990-994, 1984). Since 3-substituted- β -lapachone derivatives cause enhanced Topo I inhibition, it seems reasonable to assume that these derivatives will be effective reverse transcriptase inhibitors and prevent the replication of the HIV virus.

I claim:


1. A compound having the structure:




wherein R is $(\text{CH}_2)_n\text{-R}_1$ where n is an integer from 0-10 and R_1 is hydrogen, an alkyl, an aryl, a heteroaromatic, a heterocyclic, an alkoxy, a hydroxy, an amine, a thiol, an amide, or a halogen.

2. The compound of claim 1 wherein said compound is 3-hydroxy- β -lapachone.
3. The compound of claim 1 wherein said compound is 3-methyl- β -lapachone.
4. The compound of claim 1 wherein said compound is 3-ethoxycarbonylmethyl- β -lapachone.
5. The compound of claim 1 wherein said compound is 3-(2'-aminoethyl)- β -lapachone.
6. The compound of claim 1 wherein said compound is 3-(2'-hydroxyethyl)- β -lapachone.
7. The compound of claim 1 wherein said compound is 3-methoxy- β -lapachone.
8. The compound of claim 1 wherein said compound is 3-benzyloxy- β -lapachone.
9. The compound of claim 1 wherein said compound is 3-ethoxycarbonylmethyloxy- β -lapachone.

10. The compound of claim 1 wherein said compound is 3-allyloxy- β -lapachone.
11. A method of synthesizing a 3-substituted- β -lapachone, comprising the steps of:
 - a. alkylating a silver salt of lapachol having the formula:



to produce a mixture of a lapachol ether and a 3-substituted- β -lapachone, said 3-substituted- β -lapachone having the formula:



wherein R is $(CH_2)_n-R_1$ where n is an integer from 1-10 and R_1 is hydrogen, an alkyl, an aryl, a heteroaromatic, a heterocyclic, an

aliphatic, an alkoxy, a hydroxy, an amine, a thiol, an amide or a halogen; and

b. recovering the 3-substituted- β -lapachone.

12. The method of claim 11 further including the step of suspending the silver salt in an organic solvent prior to alkylating.

13. The method of claim 12 wherein said organic solvent is ether.

14. The method of claim 11 wherein the silver salt is alkylated in the presence of an alkyl-halide.

15. The method of claim 14 wherein said alkyl-halide is selected from the group consisting of methyl iodide, benzyl bromide, ethyl bromoacetate, 2-bromoethanol, and 2-bromoethylamine.

16. The method of claim 11 wherein recovering the 3-substituted- β -lapachone comprises the step of eluting the 3-substituted- β -lapachone from the mixture.

17. The method of claim 11 further including the step of purifying the 3-substituted- β -lapachone prior to recovering the 3-substituted- β -lapachone.

18. The method of claim 17 wherein purifying said 3-substituted β -lapachol takes place in the presence of methanol in methylene chloride.

19. The method of claim 14 wherein adding an alkyl-halide is performed while heating under reflux conditions wherein said reflux time ranges from .5 -12 hours.

20. The method of claim 12 wherein prior to suspending the silver salt of lapachol in an organic solvent, the salt is first filtered, then washed with distilled water, then washed with 95% ethanol, and then washed with ether.

21. The method of claim 14 wherein said alkyl-halide is added in a ten-fold amount relative to the silver salt.

22. The method of claim 11 wherein said 3-substituted β -lapachone is 3-allyl- β -lapachone.

23. The method of claim 11 wherein said 3-substituted β -lapachone is 3-ethoxycarbonylmethyl- β -lapachone.

24. The method of claim 11 wherein said 3-substituted β -lapachone is 3-(2'-hydroxyethyl)- β -lapachone.


25. The method of claim 11 wherein said 3-substituted β -lapachone is 3-(2'-aminoethyl)- β -lapachone.

26. The method of claim 11 wherein said 3-substituted lapachone is 3-methyl- β -lapachone.

27. The method of claim 11 wherein said 3-substituted β -lapachone is 3-hydroxy- β -lapachone.

28. A method of enhancing the lethality of a DNA damaging agent in neoplastic cells, comprising the steps of:

exposing a neoplastic cell to a 3-substituted β -lapachone having the structure:



wherein R is $(\text{CH}_2)_n\text{-R}_1$, where n is an integer from 0-10, and R_1 is hydrogen, an alkyl, an aryl, a heteroaromatic, a heterocyclic, an alkoxy, a hydroxy, an amine, a thiol, an amide, or a halogen.

29. The method of claim 28 wherein said DNA damaging agent is radiotherapy.

30. The method of claim 28 wherein said DNA damaging agent is chemotherapy.

31. The method of claim 28 wherein said 3-substituted β -lapachone is 3-ethoxycarbonylmethyl- β -lapachone.

32. The method of claim 28 wherein said 3-substituted β -lapachone is 3-(2'-Hydroxyethyl)- β -lapachone.

33. The method of claim 28 wherein said 3-substituted β -lapachone is 3-methyl- β -lapachone.

34. The method of claim 28 wherein said 3-substituted β -lapachone is 3-(2'-aminoethyl)- β -lapachone.

35. The method of claim 28 wherein said 3-substituted β -lapachone is 3-hydroxy- β -lapachone.

36. The method of claim 28 wherein said 3-substituted β -lapachone is 3-methoxy- β -lapachone.


37. The method of claim 28 wherein said 3-substituted β -lapachone is 3-benzyloxy- β -lapachone.

38. The method of claim 28 wherein said 3-substituted β -lapachone is 3-ethoxycarbonylmethoxy- β -lapachone.

39. The method of claim 28 wherein said 3-substituted- β -lapachone is 3-allyloxy- β -lapachone.

40. A method of chemotherapy comprising the steps of:

exposing a neoplastic cell to a 3-substituted β -lapachone of the formula:



wherein R is $(CH_2)_n-R_1$ where n is an integer from 0-10 and R_1 is hydrogen, an alkyl, an aryl, a heteroaromatic, a heterocyclic, an alkoxy, a hydroxy, an amine, a thiol, an amide, or a halogen in sufficient amounts to induce cell death.

41. The method of claim 40 wherein said cell death is via apoptosis.

42. The method of claim 40 wherein said cell death is via necrosis.

43. The method of claim 40 wherein said 3-substituted- β -lapachone is 3-ethoxycarbonylmethyl- β -lapachone.

44. The method of claim 40 wherein said 3-substituted- β -lapachone is 3-(2'-hydroxyethyl)- β -lapachone.

45. The method of claim 40 wherein said 3-substituted- β -lapachone is 3-methyl- β -lapachone.

46. The method of claim 40 wherein said 3-substituted β -lapachone is 3-hydroxy- β -lapachone.

47. The method of claim 40 wherein said 3-substituted- β -lapachone is 3-methoxy- β -lapachone.


48. The method of claim 40 wherein said 3-substituted- β -lapachone is 3-benzyoxy- β -lapachone.

49. The method of claim 40 wherein said 3-substituted- β -lapachone is 3-ethoxycarbonylmethoxy- β -lapachone.


50. The method of claim 40 wherein said 3-substituted- β -lapachone is 3-allyloxy- β -lapachone.

51. A method of synthesizing 3-substituted β -lapachones, comprising the steps of:

a. alkylating a 3-hydroxy- β -lapachone having the formula:



to produce a 3-O-alkyl- β -lapachone having the formula:



and;

b. recovering the 3-O-alkyl- β -lapachone.

52. The method of claim 51 further including the step of suspending the 3-hydroxy- β -lapachone in an organic solvent prior to alkylating.

53. The method of claim 52 wherein said organic solvent is dimethyl sulfoxide.

54. The method of claim 51 wherein the 3-hydroxy- β -lapachone is alkylated in the presence of an alkyl-halide.

55. The method of claim 54 wherein said alkyl-halide is selected from methyl iodide, benzyl bromide, ethyl bromoacetate, and allyl bromide.

56. The method of claim 51 wherein recovering the 3-O-alkyl- β -lapachone comprises the step of eluting the 3-O-alkyl- β -lapachone.

57. The method of claim 51 further including the step of purifying the 3-O-alkyl- β -lapachone prior to recovering the 3-O-alkyl- β -lapachone.

58. The method of claim 54 wherein alkylating in the presence of an alkyl halide is carried out at 20-25°C for about .5 hours.

59. The method of claim 51 wherein said 3-O-substituted- β -lapachone is 3-methoxy- β -lapachone.

60. The method of claim 51 wherein said 3-O-substituted- β -lapachone is 3-benzyloxy- β -lapachone.


61. The method of claim 51 wherein said 3-O-substituted- β -lapachone is 3-ethoxycarbonylmethoxy- β -lapachone.

62. The method of claim 51 wherein said 3-O-substituted- β -lapachone is 3-allyloxy- β -lapachone.

AMENDED CLAIMS

[received by the International Bureau on 17 September 1996 (17.09.96);
original claims 1-62 replaced by amended claims 1-58 (8 pages)]

1. A compound having the structure:




wherein R is allyloxy, $-(CH_2)_n-R_1$, or $-O-(CH_2)_n-R_1$, where n is an integer from 0-10 and R₁ is a hydroxy, an alkylcarboxy, a heteroaromatic, a heterocyclic, an amine, a thiol, an amide, or a halogen.


2. The compound of claim 1 wherein said compound is 3-ethoxycarbonylmethyl- β -lapachone.
3. The compound of claim 1 wherein said compound is 3-(2'-aminoethyl)- β -lapachone.
4. The compound of claim 1 wherein said compound is 3-(2'-hydroxyethyl)- β -lapachone.
5. The compound of claim 1 wherein said compound is 3-ethoxycarbonylmethyloxy- β -lapachone.
6. The compound of claim 1 wherein said compound is 3-allyloxy- β -lapachone.

7. A method of synthesizing a 3-substituted- β -lapachone, comprising the steps of:

a. alkylating a silver salt of lapachol having the formula:



to produce a mixture of a lapachol ether and a 3-substituted- β -lapachone, said 3-substituted- β -lapachone having the formula:



wherein R is hydrogen, allyloxy, $-(CH_2)_n-R_1$, or $-O-(CH_2)_n-R_1$, where n is an integer from 0-10 and R_1 is hydrogen, a hydroxy, an alkoxy, an alkylcarboxy, an aryl, a heteroaromatic, a heterocyclic, an amine, a thiol, an amide, or a halogen; and

b. recovering the 3-substituted- β -lapachone.

8. The method of claim 7 further including the step of suspending the silver salt in an organic solvent prior to alkylating.
9. The method of claim 8 wherein said organic solvent is ether.
10. The method of claim 7 wherein the silver salt is alkylated in the presence of an alkyl-halide.
11. The method of claim 10 wherein said alkyl-halide is selected from the group consisting of methyl iodide, benzyl bromide, ethyl bromoacetate, 2-bromoethanol, and 2-bromoethylamine.
12. The method of claim 7 wherein recovering the 3-substituted- β -lapachone comprises the step of eluting the 3-substituted- β -lapachone from the mixture.
13. The method of claim 7 further including the step of purifying the 3-substituted- β -lapachone prior to recovering the 3-substituted- β -lapachone.
14. The method of claim 13 wherein purifying said 3-substituted β -lapachol takes place in the presence of methanol in methylene chloride.
15. The method of claim 10 wherein adding an alkyl-halide is performed while heating under reflux conditions wherein said reflux time ranges from 0.5 to 12 hours.
16. The method of claim 8 wherein prior to suspending the silver salt of lapachol in an organic solvent, the salt is first filtered, then washed with distilled water, then washed with 95% ethanol, and then washed with ether.
17. The method of claim 10 wherein said alkyl-halide is added in a ten-fold amount relative to the silver salt.

18. The method of claim 7 wherein said 3-substituted β -lapachone is 3-allyl- β -lapachone.

19. The method of claim 7 wherein said 3-substituted β -lapachone is 3-ethoxycarbonylmethyl- β -lapachone.

20. The method of claim 7 wherein said 3-substituted β -lapachone is 3-(2'-hydroxyethyl)- β -lapachone.


21. The method of claim 7 wherein said 3-substituted β -lapachone is 3-(2'-aminoethyl)- β -lapachone.

22. The method of claim 7 wherein said 3-substituted lapachone is 3-methyl- β -lapachone.

23. The method of claim 7 wherein said 3-substituted β -lapachone is 3-hydroxy- β -lapachone.

24. A method of enhancing the lethality of a DNA damaging agent in neoplastic cells, comprising the steps of:


exposing a neoplastic cell to a 3-substituted β -lapachone having the structure:



wherein R is allyloxy, $-(CH_2)_n-R_1$, or $-O-(CH_2)_n-R_1$, where n is an integer from 0-10 and R_1 is hydrogen, a hydroxy, an alkylcarboxy, a heteroaromatic, a heterocyclic, an amine, a thiol, an amide, or a halogen.

25. The method of claim 24 wherein said DNA damaging agent is radiotherapy.
26. The method of claim 24 wherein said DNA damaging agent is chemotherapy.
27. The method of claim 24 wherein said 3-substituted β -lapachone is 3-ethoxycarbonylmethyl- β -lapachone.
28. The method of claim 24 wherein said 3-substituted β -lapachone is 3-(2'-hydroxyethyl)- β -lapachone.
29. The method of claim 24 wherein said 3-substituted β -lapachone is 3-methyl- β -lapachone.
30. The method of claim 24 wherein said 3-substituted β -lapachone is 3-(2'-aminoethyl)- β -lapachone.
31. The method of claim 24 wherein said 3-substituted β -lapachone is 3-hydroxy- β -lapachone.
32. The method of claim 24 wherein said 3-substituted β -lapachone is 3-methoxy- β -lapachone.
33. The method of claim 24 wherein said 3-substituted β -lapachone is 3-benzyloxy- β -lapachone.
34. The method of claim 24 wherein said 3-substituted β -lapachone is 3-ethoxycarbonylmethoxy- β -lapachone.
35. The method of claim 28 wherein said 3-substituted- β -lapachone is 3-allyloxy- β -lapachone.

36. A method of chemotherapy comprising the steps of:
exposing a neoplastic cell to a 3-substituted β -lapachone of the formula:



- wherein R is allyloxy, $-(CH_2)_n-R_1$, or $-O-(CH_2)_n-R_1$, where n is an integer from 0-10 and R_1 is hydrogen, a hydroxy, an alkylcarboxy, a heteroaromatic, a heterocyclic, an amine, a thiol, an amide, or a halogen, in sufficient amounts to induce cell death.
37. The method of claim 36 wherein said cell death is via apoptosis.
38. The method of claim 36 wherein said cell death is via necrosis.
39. The method of claim 36 wherein said 3-substituted- β -lapachone is 3-ethoxycarbonylmethyl- β -lapachone.
40. The method of claim 36 wherein said 3-substituted- β -lapachone is 3-(2'-hydroxyethyl)- β -lapachone.
41. The method of claim 36 wherein said 3-substituted- β -lapachone is 3-methyl- β -lapachone.
42. The method of claim 36 wherein said 3-substituted β -lapachone is 3-hydroxy- β -lapachone.
43. The method of claim 36 wherein said 3-substituted- β -lapachone is 3-methoxy- β -lapachone.


44. The method of claim 36 wherein said 3-substituted- β -lapachon is 3-benzyloxy- β -lapachone.

45. The method of claim 36 wherein said 3-substituted- β -lapachone is 3-ethoxycarbonylmethoxy- β -lapachone.


46. The method of claim 36 wherein said 3-substituted- β -lapachone is 3-allyloxy- β -lapachone.

47. A method of synthesizing 3-substituted β -lapachones, comprising the steps of:

a. alkylating a 3-hydroxy- β -lapachone having the formula:



to produce a 3-O-alkyl- β -lapachone having the formula:



and;

b. recovering the 3-O-alkyl- β -lapachone.

48. The method of claim 47 further including the step of suspending the 3-hydroxy- β -lapachone in an organic solvent prior to alkylating.

49. The method of claim 48 wherein said organic solvent is dimethyl sulfoxide.

50. The method of claim 47 wherein the 3-hydroxy- β -lapachone is alkylated in the presence of an alkyl-halide.

51. The method of claim 50 wherein said alkyl-halide is selected from methyl iodide, benzyl bromide, ethyl bromoacetate, and allyl bromide.

52. The method of claim 47 wherein recovering the 3-O-alkyl- β -lapachone comprises the step of eluting the 3-O-alkyl- β -lapachone.

53. The method of claim 47 further including the step of purifying the 3-O-alkyl- β -lapachone prior to recovering the 3-O-alkyl- β -lapachone.

54. The method of claim 50 wherein alkylating in the presence of an alkyl halide is carried out at 20-25°C for about 0.5 hours.

55. The method of claim 47 wherein said 3-O-substituted- β -lapachone is 3-methoxy- β -lapachone.

56. The method of claim 47 wherein said 3-O-substituted- β -lapachone is 3-benzyloxy- β -lapachone.

57. The method of claim 47 wherein said 3-O-substituted- β -lapachone is 3-ethoxycarbonylmethoxy- β -lapachone.

58. The method of claim 47 wherein said 3-O-substituted- β -lapachone is 3-allyloxy- β -lapachone.




Fig. 1

Apoptosis Assay Using HL60 Cells




Fig. 1a




Fig. 2




FIG. 3

5/7




Fig. 4

6/7




FIG. 5




Fig. 6

INTERNATIONAL SEARCH REPORT

Intern. Appl. No.

PCT/US 96/05673

A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C07D311/92 A61K31/35

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D A61K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO,A.94 04145 (DANA FARBER INSTIT.) 3 March 1994 see the whole document	1,3,6,7, 28,40
X	--- CHEMICAL ABSTRACTS, vol. 110, no. 1, 1989 Columbus, Ohio, US; abstract no. 88167h, D.BOOTHMAN ET AL.: "INHIBITION OF POTENTIALLY LETAL DNA DAMAGE REPAIR IN HUMAN TUMOR CELLS BY BETA-LAPACHONE" page 35; XP002007963 see abstract	1,28, 40-42
X	& CANCER RES., vol. 49, no. 3, 1989, ENGL., pages 605-612, --- -/--	1,28, 40-42

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

9 July 1996

Date of mailing of the international search report

18.07.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Francois, J

INTERNATIONAL SEARCH REPORT

Intern. Application No

PCT/US 96/05673

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CHEMICAL ABSTRACTS, vol. 91, no. 1, 1979 Columbus, Ohio, US; abstract no. 151101p, R.DOCAMPO ET AL.: "BETA-LAPACHONE ENHANCEMENT OF LIPID PEROXYDATION" page 25; XP002007964 see abstract	1,28, 40-42
X	& BIOCHEM. PHARMACOL., vol. 28, no. 6, 1979, ENGL, pages 723-728, ---	1,28, 40-42
X	CHEMICAL ABSTRACTS, vol. 112, no. 1, 1990 Columbus, Ohio, US; abstract no. 30301x, D.BOOTHMAN: "CAN A DNA REPAIR INHIBITOR BLOCK THE TUMORIGENIC TRANSFORMATION OF NORMAL CELLS FOLLOWING A GENETIC INSULT?" page 19; XP002007965 see abstract	1,28, 40-42
X	& BIOL. ZENTRALBLAT., vol. 108, no. 5, 1989, ENGL., pages 415-421, ---	1,28, 40-42
X	CHEMICAL ABSTRACTS, vol. 122, no. 25, 1995 Columbus, Ohio, US; abstract no. 309888w, I.SZUMIEL ET AL.: "EFFECTS OF TOPOISOMERASE I-TARGETED DRUGS ON RADIATION RESPONSE OF L5178Y" page 460; XP002007966 see abstract	1,28, 40-42
X	& INT. J. RADIAT, BIOL., vol. 67, no. 4, 1995, ENGL., pages 441-448, ---	1,28, 40-42
X	CHEMICAL ABSTRACTS, vol. 123, no. 1, 1995 Columbus, Ohio, US; abstract no. 217883h, S.PLANCHON ET AL.: "BETA-LAPACHONE MEDIATED APOPTOSIS IN HUMAN PROMYELOCYTIC LEUKEMIA(HL-60) AND HUMAN PROSTATE CANCER CELLS" page 43; XP002007967 see abstract	1,28, 40-42
X	& CANCER RES., vol. 55, no. 17, 1995, ENGL., pages 3706-3711, ---	1,28, 40-42

-/--

INTERNATIONAL SEARCH REPORT

Intern. Application No

PCT/US 96/05673

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	CHEMICAL ABSTRACTS, vol. 123, no. 1, 1995 Columbus, Ohio, US; abstract no. 217884j, CHIANG LI ET AL.: "INDUCTION OF APOPTOSIS BY BETA-LAPACHONE IN HUMAN PROSTATE CANCER CELLS." page 43; XP002007968 see abstract	1,28, 40-42
X	& CANCER RES., vol. 55, no. 17, 1995, ENGL., pages 3712-3715, ---	1,28, 40-42
X	PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES OF USA, vol. 86, no. 13, July 1989, WASHINGTON US, pages 4963-4967, XP002007962 D.BOOTHMAN ET AL.: "INHIBITION OF RADIATION INDUCED NEOPLASTIC TRANSFORMATION BY BETA-LAPACHONE" see page 4963 - page 4966 ---	1,28, 40-42
X	CHEMICAL ABSTRACTS, vol. 78, no. 27, 1973 Columbus, Ohio, US; abstract no. 147733k, J.MOCK ET AL.: "CHEM. STUDIES OF THE PROTEACEAE VI." page 361; XP002007969 see abstract	1,2
X	& AUST. J. CHEM., vol. 26, no. 5, 1973, SYDNEY, pages 1121-1130, ---	1,2
X	CHEMICAL ABSTRACTS, vol. 87, no. 17, 1977 Columbus, Ohio, US; abstract no. 127264r, A.PINTO ET AL.: "SCHSTOSOMA MANSONIBLOCKAGE OF CERCARIAL SKIN PENETRATION BY CHEMICAL AGENTS. I." page 58; XP002007970 see abstract	1,2,28
X	& TRANS R. SOC. TROP. MED. HYG., vol. 71, no. 2, 1977, BRAZIL, pages 133-135, -----	1,2,28

INTERNATIONAL SEARCH REPORT

Information on patent family members

Intern. Appl. Application No

PCT/US 96/05673

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9404145	03-03-94	NONE	